

# **Analysis of CO Emission in Comet C/2002 T7 (LINEAR) from Infrared Observations**

**William M. Anderson Jr.**

*Department of Physics  
Catholic University of America  
Washington, DC 20064*

*and*

*Solar System Exploration Division  
NASA Goddard Space Flight Center  
Greenbelt, MD 20711*

[88anderw@cua.edu](mailto:88anderw@cua.edu)

**Michael A. DiSanti**

*Goddard Center for Astrobiology  
and*

*Solar System Exploration Division  
NASA Goddard Space Flight Center  
Greenbelt, MD 20711*

**Michael J. Mumma**

*Goddard Center for Astrobiology  
and*

*Solar System Exploration Division  
NASA Goddard Space Flight Center  
Greenbelt, MD 20711*

**Neil Dello Russo**

*Department of Physics  
Catholic University of America  
Washington, DC 20064*

*and*

*Solar System Exploration Division  
NASA Goddard Space Flight Center  
Greenbelt, MD 20711*

**Boncho P. Bonev**

*Department of Physics  
University of Toledo  
Toledo, OH 43066*

*and*

*Solar System Exploration Division  
NASA Goddard Space Flight Center  
Greenbelt, MD 20711*

**Karen Magee-Sauer**

*Department of Physics & Astronomy  
Rowan University  
Glassboro, NJ 08028*

**Erika L. Gibb**

*Department of Physics  
University of Notre Dame*

Cometary nuclei are the most primitive remnants of the early Solar System. Their physical and chemical attributes allow a glimpse into the conditions in which icy bodies formed. Only in recent years has it been possible to routinely study parent volatiles in the infrared. A significant variation in composition among ten comets sampled to date has been demonstrated, and this forms the foundation of a new cometary taxonomy based on chemistry.

In spring 2004, we observed comet C/2002 T7 (LINEAR) using the facility echelle spectrometer (CSHELL) at the NASA Infrared Telescope Facility on Mauna Kea, Hawaii. CSHELL offers seeing-limited spatial resolution and sufficiently high spectral resolving power ( $R \sim 2.5 \times 10^4$ ) to permit line-by-line intensities to be measured along its 30 arcsecond-long slit. Emission lines from multiple molecular species were targeted in the 3 to 5 micron infrared region, and our observations revealed an extremely rich chemistry in comet T7. Here we present production rates, mixing ratios, and rotational temperatures for CO spanning UT 3 - 9 May 2004, based on preliminary analysis of lines in the R and P branches of the  $v = 1 - 0$  fundamental ro-vibrational band near 4.7 microns. Through comparison with abundances of other oxygen-bearing molecules, specifically formaldehyde ( $\text{H}_2\text{CO}$ ) and methyl alcohol ( $\text{CH}_3\text{OH}$ ), potential implications for the comet's volatile carbon-oxygen history will be discussed. The prospective of laboratory ice irradiation experimentation, as a comparative method to cometary observations, will also be explored.

This research is supported by the NASA Planetary Astronomy Program, through RTOP 344-32-98 and 344-32-51 to M. A. DiSanti.